

Open Access Original Article



Coronary external diameter index for assessing coronary artery involvement in Kawasaki disease

Andrea Azzarelli^{1*}, Cuono Cucco², Lauro Cortigiani¹, Marco Scalese³, Valentina Annoni⁴, Francesco Bovenzi¹, Raffaele Domenici¹, Angelina Vaccaro¹, Francesco Vierucci¹

¹Cardiology Division, San Luca Hospital, 55100 Lucca, Italy

²Cardiology Division, University Hospital Cisanello, 56120 Pisa, Italy

³Institute of Clinical Physiology, National Research Council, 56100 Pisa, Italy

⁴Department of Medicine, Versilia Hospital, 55041 Lucca, Italy

***Correspondence**: Andrea Azzarelli, Cardiology Division, San Luca Hospital, Via Francesconi 556, 55100 Lucca, Italy. aazzarelli60@gmail.com

Academic Editor: Erkin Mirrakhimov, Kyrgyz State Medical Academy, Kyrgyzstan Received: January 21, 2023 Accepted: May 6, 2023 Published: October 31, 2023

Cite this article: Azzarelli A, Cucco C, Cortigiani L, Scalese M, Annoni V, Bovenzi F, et al. Coronary external diameter index for assessing coronary artery involvement in Kawasaki disease. Explor Cardiol. 2023;1:103–13. https://doi.org/10.37349/ec. 2023.00011

Abstract

Aim: Transthoracic echocardiography is commonly used to assess coronary artery dilatation in Kawasaki disease (KD). However, existing criteria often miss early abnormalities. This study examines the utility of a new parameter, coronary external diameter index (CEDi), for early diagnosis and monitoring in KD.

Methods: CEDi of left main (LM) and right coronary artery (RCA), calculated as the ratio of coronary artery external diameter (i.e., the distance between the outer coronary edges measured in the proximal segment of the artery) and the diameter of the aortic annulus, was evaluated in 34 patients (age 23 mouths ± 13 months) with KD at the hospital admission and after 2 weeks and 8 weeks of treatment. The control group consisted of 210 healthy children aged 20 months ± 13.4 months. Z-score charts for LM and RCA coronary external diameter (CED) were obtained.

Results: Compared with controls, KD patients had a markedly higher mean value of LM CEDi (0.53 ± 0.06 vs. 0.33 ± 0.04 ; P < 0.0001) and RCA CEDi (0.48 ± 0.05 vs. 0.31 ± 0.04 ; P < 0.0001) at hospital admission. By ROC analysis, LM CEDi of 0.41, and RCA coronary artery thickness index (CATi) of 0.39 were the best cutoffs to confirm the clinical diagnosis of KD, both exhibiting 100% sensitivity and specificity. Mean LM CEDi and RCA CEDi values decreased significantly (P < 0.0001) after 2 weeks of follow-up and were similar to controls (P = 0.53 and P = 0.12, respectively) 8 weeks after admission.

Conclusions: In patients with KD, CEDi of LM and RCA is an accurate parameter to evaluate coronary artery involvement in the early phase of the illness and during follow-up.

Keywords

Kawasaki disease, echocardiography, Z-score, coronary artery dilatation

© The Author(s) 2023. This is an Open Access article licensed under a Creative Commons Attribution 4.0 International License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, sharing, adaptation, distribution and reproduction in any medium or format, for any purpose, even commercially, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.



Introduction

Coronary artery aneurysm is the most serious complication of Kawasaki disease (KD), with prevalence ranging from 25% in untreated patients to 8% in those treated with a four low-dose immunoglobulin protocol to 5% in those treated with a single high-dose immunoglobulin protocol [1-5]. Transthoracic echocardiography is the first-line imaging technique to evaluate coronary lumen dilatation and aneurysm growth in KD [1-3, 6-8], allowing the classification of coronary damage [9, 10]. In particular, the coronary artery Z-score represents the widely used method to assess coronary artery dilatation in KD [11-17]. However, it shows a normal value in about 75% of cases at the time of hospital admission as the dilatation usually develops 7–10 days later [6-10]. Therefore, an early normal echocardiogram does not rule out the diagnosis of KD [6, 18], which can cause severe clinical relapses particularly in subjects with clinically incomplete clinical presentation [18, 19].

In this prospective study, we investigated the usefulness of a new echocardiographic parameter, such as the external coronary external diameter index (CEDi), for the early identification and follow-up of coronary artery involvement in patients with KD.

Materials and methods

Patients

From January 2010 to September 2020, 42 patients with KD were admitted to the Pediatric Department of San Luca Hospital, Lucca, Italy, and underwent transthoracic echocardiography. Of these patients, 2 were admitted from March 2020 to September 2020 during the coronavirus disease (COVID) pandemic. Two patients were excluded due to an uncooperative attitude. Of the remaining 40 patients, 6 (15%) were excluded due to inadequate imaging of left main (LM; n = 1) or right coronary artery (RCA) lumen (n = 1) or coronary artery thickness (CAT; n = 4). Thus, 34 patients, including 17 males; mean [± SD (standard deviation)] age 23 months ± 13 months, formed the study group. Of them, 28 met the diagnostic criteria for KD [1, 6, 12], while 6 patients had an incomplete clinical presentation of KD, and the diagnosis was confirmed by additional abnormal laboratory criteria.

Complete clinical recovery was achieved in all patients after treatment with aspirin and a single (n = 31) or double infusion of high-dose (2 g/kg) immunoglobulin (n = 3).

The control group consisted of consecutive children aged between 2 months and 48 months and with echocardiographic evidence of structurally and functionally normal heart who were referred from January 2010 to December 2013 to the outpatient echo lab of the San Luca Hospital for evaluation of heart murmur, musculoskeletal pain, electrocardiographic abnormalities, family history of congenital heart defects, minimal and hemodynamically insignificant findings such as patent foramen ovale, mild tricuspid or pulmonary valve regurgitation. Of the 265 initial children, 34 were excluded due to uncooperative attitude. Of the remaining 231 children, 21 (9%) were excluded due to inadequate imaging of the coronary artery lumen (3 LM, 3 RCA, 1 both arteries) or CAT (10 LM, 3 RCA, 1 both arteries). Consequently, the remaining 210 healthy children [125 males; mean (± SD) age 20 months ± 14 months] served as controls. The enrollment into the study is shown in Figure 1, reporting how many individuals were excluded at each exclusion step.

Coronary artery measurements

Echocardiographic studies were performed using a Philips iE33 Ultrasound System equipped with a highfrequency phased-array sector scanning probe (S8-3 ultrasound probe by Philips Healthcare) and second harmonic technology. The aortic annulus diameter, coronary lumen diameter, coronary external diameter (CED), CED Z-score, CEDi, and CAT of LM and RCA were evaluated in controls and KD patients at hospital admission. In these last subjects the same parameters were assessed also after 2 weeks and 8 weeks of treatment.



Figure 1. Consort diagram showing how many individuals were excluded at each exclusion step

The diameter of the aortic annulus was measured in the standard parasternal long axis view, at early systole, inner to inner edge from hinge point to hinge point of the aortic cusps (Figure 2) [16].



Figure 2. Parasternal long axis view showing measurement of the aortic annulus during end-systole

The short-axis parasternal view was used for LM and RCA measurements. In particular, the coronary lumen was obtained as the distance between the inner edges of the arteries [6, 13, 14], while CED as the distance between the outer edges of the arteries, thus including the lumen diameter and the thickness of the arterial wall (Figure 3). The measurements were made during early diastole and the optimal point was chosen 3–5 mm from the origin of the coronary artery. In the case of inadequate imaging of the coronary arteries from a short-axis parasternal view, a modified 5-chamber apical approach was adopted (Figure 4). At least 5 measurements were made in each subject and the results were averaged. All measurements were normalized by body surface area (BSA), calculated using the Haycock formula [BSA = $0.024265 \times \text{height}$ (cm)^{0.3964} × weight (kg)^{0.5378}], which is the best predictor of cardiovascular sizes [11, 12].

LM and RCA CEDi were obtained from the ratio of the external diameter of the coronary artery to the diameter of the aortic annulus. Examples of the calculation of LM and RCA CEDi in a normal subject and a patient with KD are shown in Figure 5.

LM coronary external diameter

LM coronary esternal diameter

RCA coronary external diameter

RCA coronary external diameter



Figure 3. Parasternal short axis view showing LM (left) and right (right) CED measurement as the distance between the outer edges of the arteries



Figure 4. Apical modified 5-chamber view showing LM (left) and right (right) CED measurement as the distance between the outer edges of the arteries

CAT was obtained as the difference between CED and internal coronary lumen. All the studies were performed by the same senior sonographer (Andrea Azzarelli).

Statistical analysis

Continuous variables are expressed as mean \pm SD. Differences between groups were compared using Student's *t* and chi-square test, as appropriate. Correlations of CED and CEDi with age, weight, height, and BSA in the control group were estimated with Pearson's coefficients. A receiver operating characteristic analysis was adopted to determine the best value of LM CEDi and RCA CEDi to confirm the clinical diagnosis of KD. Statistical significance was set at *P* < 0.05. Inter-observer and intra-observer agreement was calculated based on overall agreement (observed percent exact agreement) and tested using linear regression analysis.

By analyzing the data of the 210 subjects of the control group, the Z-score tables for LM and RCA CAT were obtained. Since the assumption of logistic regression is that the residuals should be normally distributed, the Shapiro-Wilk and Kolmogorov-Smirnov tests have been used to evaluate the normality of the residuals [11]. The regression equation with the highest R^2 (goodness-of-fit) was selected for LM and RCA CAT. The heteroskedasticity used to describe the behavior of the variance of the residuals was then verified using the White test and the Breusch-Pagan test. The statistical package for the social sciences (IBM SPSS Statistics V22.0) was used for the analysis.



Figure 5. Comparative analysis of LM and RCA CEDi calculations in a patient with KD and a healthy subject. The left panels show the calculation of LM and RCA CEDi in a patient with KD. The aortic annulus (top), LM CED (middle), and RCA CED (bottom) were 12.4 mm, 7.4 mm, and 7.9 mm, respectively. Thus, LM CEDi of 0.60 and RCA CEDi of 0.63 were derived. The right panels show the calculation of LM and RCA CEDi in a normal subject. The aortic annulus (top), LM CED (middle), and RCA CED (bottom) were 12.1 mm, 3.2 mm, and 2.4 mm, respectively. Thus, LM CEDi of 0.26 and RCA CEDi of 0.20 were derived

Results

Out of the total population included in the study (KD patients and controls), the number of subjects who were able to undergo an adequate examination of both coronary arteries was 244/307 (79%).

The following Table 1 shows the best-fit model, its R^2 value, and the regression equation that allows deriving the Z-score plots for LM and RCA CED (Figure 6) in the group of 210 normal subjects. In these individuals, LM and RCA CED correlated significantly with age, height, weight, and BSA, whereas LM and RCA CED i failed to do it (Table 2).

|--|

Measurement	Intercept	В	SEE (√MSE)	R ²	SW	KS	BP	W
RCA CED	1.506	0.448	0.128	0.484	0.255	0.087	0.471	0.600
LM CED	1.605	0.471	0.134	0.485	0.385	0.200	0.114	0.187
BSA Haycock. (In[y] = a + b*ln[x]); Z value = exp((In[Measurement] – (Intercept + B*ln[BSA]))/\ImegABE)								

B: regression coefficient; SEE (\sqrt{MSE}): standard error of the estimate; R^2 : coefficient of determination; SW: Shapiro-Wilk test; KS: Kolmogorov-Smirnov test; BP: Breusch-Pagan test; W: Wald test



Figure 6. Percentile charts for LM CED (top), and RCA CED (bottom) according to BSA calculated using the Haycock formula

CED and index	Age		Height	Height		Weight		BSA	
	r	P-value	r	P-value	r	P-value	r	P-value	
LM CED	0.63	< 0.001	0.66	< 0.001	0.68	< 0.001	0.68	< 0.001	
RCA CED	0.38	< 0.001	0.43	< 0.001	0.46	< 0.001	0.46	< 0.001	
LM CEDi	0.08	0.22	0.04	0.51	0.06	0.35	0.06	0.39	
RCA CEDi	0.02	0.74	0.13	0.90	0.01	0.84	0.01	0.90	

The main clinical and echocardiographic parameters in KD patients on admission and in controls are shown in Table 3. No difference between groups was observed for age, height, weight, and BSA, while aortic annulus diameter was greater (P = 0.001) in subjects with KD.

Table 3. Clinical and echocardiographic characteristics of patients with KD at admission and controls

Clinical and echocardiographic characteristics	KD patients (<i>n</i> = 34)	Control subjects (<i>n</i> = 210)	P-value	
Age (months)	23 ± 13	20 ± 14	0.15	
Height (cm)	84 ± 14	81 ± 16	0.28	
Weight (kg)	11 ± 3	11 ± 4	0.63	
BSA	0.51 ± 0.10	0.49 ± 0.13	0.44	
Aortic annulus diameter (mm)	11.8 ± 1.5	10.7 ± 1.7	0.001	
LM lumen diameter (mm)	2.3 ± 0.7	1.6 ± 0.3	< 0.0001	
LM lumen Z-score	0.5 ± 0.3	0.2 ± 0.1	< 0.0001	
LM lumen Z-score ≥ 2	4 (12%)	0	< 0.0001	
LM CED (mm)	6.3 ± 1.1	3.6 ± 0.6	< 0.0001	
LM CED Z-score	4.1 ± 1.2	0.01 ± 1	< 0.0001	
LM CAT (mm)	3.9 ± 0.9	1.9 ± 0.5	< 0.0001	

Table 3. Clinical and echocardiographic characteristics of patients with KD at admission and controls (continued)

Clinical and echocardiographic characteristics	KD patients (<i>n</i> = 34)	Control subjects (<i>n</i> = 210)	P-value
LM CEDi	0.5 ± 0.1	0.3 ± 0.1	< 0.0001
RCA lumen diameter (mm)	2.0 ± 0.5	1.6 ± 0.3	< 0.0001
RCA lumen Z-score	0.4 ± 0.2	0.2 ± 0.1	< 0.0001
RCA lumen Z-score ≥ 2	3 (9%)	0	< 0.0001
RCA CED (mm)	5.7 ± 1.1	3.3 ± 0.6	< 0.0001
RCA CED Z-score	4.1 ± 1.2	0.01 ± 0.9	< 0.0001
RCA CAT (mm)	3.7 ± 1.0	1.7 ± 0.4	< 0.0001
RCA CEDi	0.4 ± 0.1	0.3 ± 0.1	< 0.0001

CED, CEDi and CAT of LM and RCA were markedly higher (P < 0.0001) in KD patients than in controls (Table 3). The former had also a larger lumen diameter (P < 0.0001) of either LM or RCA. However, an abnormal Z-score (≥ 2) for LM and RCA at admission was found in only 4 (12%) and 3 (9%) KD patients, respectively (Table 3).

With a ROC analysis, LM CEDi of 0.41, and RCA CEDi of 0.39 at hospital admission were the best cut-offs to confirm the clinical diagnosis of KD, both showing 100% sensitivity and specificity.

In KD patients, CEDi of LM and RCA decreased progressively at 2 weeks and 8 weeks of follow-up (Figure 7). In particular, at 8 weeks the mean value of LM and RCA CEDi were comparable to those of controls (P = 0.53 and P = 0.12, respectively), and 33 of 34 patients had LM CEDi < 0.41 and RCA CEDi <0.39. The mean values of CED, CED Z-score, CEDi, and CAT of LM and RCA in KD patients at admission, and 2 weeks and 8 weeks after hospitalization, as shown in Table 4.



Figure 7. Behavior of LM (left) and RCA CEDi (right) values evaluated at hospital admission, after 2 weeks and 8 weeks in the 34 patients with KD

 Table 4.
 Values of CED, CED Z-score, CEDi, and CAT of LM and RCA in KD patients at admission, and 2 weeks and 8 weeks after hospitalization

Echo Stages	LM CED	LM CED Z-score	LM CEDi	LM CAT	RCA CED	RCA CED Z-score	RCA CEDi	RCA CAD
Admission	6.3 ± 1.1	4.1 ± 1.2	0.5 ± 0.1	3.9 ± 0.9	5.7 ± 1.1	4.1 ± 1.2	0.4 ± 0.1	3.7 ± 1.0
2 weeks	4.2 ± 0.8	0.8 ± 1.2	0.4 ± 0.1	2.6 ± 0.7	3.9 ± 0.1	0.8 ± 1.1	0.3 ± 0.1	2.3 ± 0.8
8 weeks	4.0 ± 0.5	0.5 ± 0.7	0.3 ± 0.1	2.1 ± 0.7	3.5 ± 1.1	0.3 ± 0.9	0.3 ± 0.1	1.9 ± 0.7

In the assessment of inter-observer variability by 2 independent observers (Andrea Azzarelli, Lauro Cortigiani Cuono Cucco) in a set of 40 individuals (20 with KD and 20 controls), a significant correlation

was found for LM CED (r = 0.97), RCA CED (r = 0.96), LM CEDi (r = 0.96), and RCA CEDi (r = 0.95), and the agreement for LM CEDi > or < 0.41 and RCA CEDi > or < 0.39 was evidenced in all cases. In the same way, the intra-observer variability was very high (r = 0.99) for both CED and CEDi.

Discussion

In this prospective study on KD patients with a relatively low level of coronary dilatation, we found that CEDi of LM and RCA is a feasible and accurate tool for assessing and monitoring coronary artery involvement. In particular, LM CEDi \geq 0.41 and RCA CEDi \geq 0.39 at hospital admission provided a diagnostic sensitivity and specificity of 100%, allowing confirmation of the diagnosis of KD even in the case of incomplete clinical presentation. Of interest, the parameter progressively decreased to normalize in 97% of cases at 8 weeks of follow-up.

CEDi shows a stable value in the whole age group considered as it is not affected by BSA. However, the aortic annulus diameter is the determinant for its calculation, and CEDi cannot be used in children with aortic disease accordingly. For this reason, we provided Z-score nomogram charts describing the correlation of LM and RCA with BSA. Our charts are practical and easy to applicate clinicians. In fact, by measuring the external coronary artery diameter and using the Z-score normograms it is possible to determine the normal range regardless of the presence of aortic artery disease.

Comparison with previous studies

KD is characterized by systemic vasculitis involving the medium-sized arteries of multiple organs including the heart [6, 20–22]. Coronary artery involvement can result in a coronary artery aneurysm, ranging from a slight dilatation to the formation of a giant aneurysm [1, 7, 23, 24], usually appearing eight days after the onset of fever. Although the incidence of coronary artery aneurysm has been reduced to approximately 5% by immunoglobulin treatment [1-4, 9, 10, 23], it remains the most serious complication of KD. Echocardiography is the first-line method for identifying and characterizing changes in the coronary arteries, due to its non-invasive nature and good accuracy. Of note, the coronary artery lumen can be evaluated in most segments of the coronary artery [1–3, 5, 25–27]. The first classification of coronary dilatation was based on the absolute or relative lumen diameter, regardless of patient size, weight, or BSA [24]. In recent years the Z-score method, which normalizes the lumen of the coronary artery to the BSA, representing the number of standard deviations from the mean, has been increasingly adopted to evaluate coronary artery dilatation [12, 17]. A classification of coronary dilatation based on Z-score and raw size has recently been recommended [6]. However, approximately 75% of KD patients have normal coronary arteries on admission echocardiogram, and 30% to 50% of those with a normal coronary lumen (Z-score < 2) show a progressive reduction in lumen size during follow-up, suggesting initial dilatation that resolves within 4–8 weeks [6–9]. Consequently, normal or mild dilatation of the coronary lumen has a low positive predictive value for the early diagnosis of KD, adversely affecting the accuracy of the echocardiogram at admission [28–30]. This lack of sensitivity of the echocardiographic technique is particularly relevant in individuals with clinically incomplete KD and in children less than 1 year of age who are at increased risk of coronary aneurysms [18, 19]. Our results confirm the lack of sensitivity of the LM and RCA lumen Z-score at hospital admission, demonstrating abnormal values in only 12% and 9% of cases, respectively. Conversely, LM CEDi \ge 0.41 and RCA CEDi \ge 0.39 were observed in all patients with KD at hospital admission and in no subjects of the control group.

Clinical implications

The increase in CEDi in KD is likely due to the involvement of the coronary wall and pericoronary tissue during the acute inflammatory phase, thus resulting not only in the slight dilatation of the coronary lumen but also in the increased thickness of the coronary artery wall. Previous reports with ultrasound and nonultrasound (optical coherence tomography) techniques have shown the increased coronary wall thickness in the acute phase of KD, and its reversibility with treatment [20, 31, 32], corroborating the results of the current study reporting markedly higher mean values of CAT at admission in KD patients compared to controls.

Assessment of CEDi may have important clinical implications for the management of patients with KD. Since CEDi of LM and RCA was increased in all of our patients at hospital admission but in no control, it may represent a useful additional parameter to confirm the diagnosis of KD allowing for early starting medical therapy. Furthermore, the serial evaluation of CEDi during the follow-up can allow use to evaluate the evolution of the coronary involvement.

Study limitations

Our study is single center and all examinations were performed by the same expert pediatric echocardiographer. Furthermore, the number of patients is quite small despite the 10-year enrollment period, which is largely due to the relatively low incidence of KD in our country. Therefore, it would be desirable if our preliminary results were confirmed by a multicenter study conducted on a larger patient population. Sedation for uncooperative patients was not available in our echo lab and this fact did not allow use to investigate all consecutive patients hospitalized for KD thus reducing the feasibility of the study. Since the control group consisted of afebrile subjects without echocardiographic evidence of cardiac disease, it remains to be defined whether CEDi is a specific marker of KD. The size of the diameter of the aortic annulus was slightly greater in KD patients than in controls confirming the results of previous experiences [29, 30]. However, this fact seems to strengthen our findings by lowering the CEDi value in KD patients compared to controls.

Abbreviations

BSA: body surface area CAT: coronary artery thickness CED: coronary external diameter CEDi: coronary external diameter index KD: Kawasaki disease LM: left main RCA: right coronary artery

Declarations

Author contributions

AA, CC and LC: Conceptualization, Investigation, Writing—original draft. VA, FB, RD, AV and FV: Investigation, Date curation. MS and CC: Formal Analysis. All authors read and approved the submitted version.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Ethical approval

The study complies with the Declaration of Helsinki and was approved by the institutional ethic committee (Azienda USL Toscana Nordovest, decreto N. 1954, June 19, 2018).

Consent to participate

Informed consent to participate in the study was obtained from all participants.

Consent to publication

Informed consent to publication was obtained from relevant participants.

Availability of data and materials

Not applicable.

Funding

Not applicable.

Copyright

© The Author(s) 2023.

References

- 1. Marchesi A, Tarissi de Jacobis I, Rigante D, Rimini A, Malorni W, Corsello G, et al. Kawasaki disease: guidelines of the Italian Society of Pediatrics, part I – definition, epidemiology, etiopathogenesis, clinical expression and management of the acute phase. Ital J Pediatr. 2018;44:102.
- 2. Dajani AS, Taubert KA, Gerber MA, Shulman ST, Ferrieri P, Freed M, et al. Diagnosis and therapy of Kawasaki disease in children. Circulation. 1993;87:1776–80.
- 3. Altman CA. Clinical assessment of coronary arteries in Kawasaki disease: focus on echocardiographic assessment. Congenit Heart Dis. 2017;12:636–40.
- 4. Suzuki A, Tizard EJ, Gooch V, Dillon MJ, Haworth SG. Kawasaki disease: echocardiographic features in 91 cases presenting in the United Kingdom. Arch Dis Child. 1990;65:1142–6.
- 5. Margossian R, Lu M, Minich LL, Bradley TJ, Cohen MS, Li JS, et al. Predictors of coronary artery visualization in Kawasaki disease. J Am Soc Echocardiogr. 2011;24:53–9.
- 6. McCrindle BW, Rowley AH, Newburger JW, Burns JC, Bolger AF, Gewitz M, et al. Diagnosis, treatment, and long-term management of Kawasaki disease: a scientific statement for health professionals from the American Heart Association. Circulation. 2017;135:e927–99.
- 7. McCrindle BW, Li JS, Minich LL, Colan SD, Atz AM, Takahashi M, et al. Coronary artery involvement in children with Kawasaki disease: risk factors from analysis of serial normalized measurements. Circulation. 2007;116:174–9.
- 8. Kim SH, Kim JY, Kim GB, Yu JJ, Choi JW. Diagnosis of coronary artery abnormalities in patients with Kawasaki disease according to established guidelines and z score formulas. J Am Soc Echocardiogr. 2021;34:662–72.e3.
- 9. Tulloh RM, Wood LE. Coronary artery changes in patients with Kawasaki disease. Acta Paediatr. 2004; 93:75–9.
- 10. Crystal MA, Manlhiot C, Yeung RS, Smallhorn JF, McCrindle BW. Coronary artery dilation after Kawasaki disease for children within the normal range. Int J Cardiol. 2009;136:27–32.
- 11. Cantinotti M, Scalese M, Murzi B, Assanta N, Spadoni I, Festa P, et al. Echocardiographic nomograms for ventricular, valvular and arterial dimensions in Caucasian children with a special focus on neonates, infants and toddlers. J Am Soc Echocardiogr. 2014;27:179–91.e2.
- 12. Lopez L, Colan S, Stylianos M, Granger S, Trachtenberg F, Frommelt P, et al.; Pediatric Heart Network Investigators. Relationship of echocardiographic Z scores adjusted for body surface area to age, sex, race, and ethnicity: the pediatric heart network normal echocardiogram database. Circ Cardiovasc Imaging. 2017;10:e006979.
- 13. Dallaire F, Dahdah N. New equations and a critical appraisal of coronary artery Z scores in healthy children. J Am Soc Echocardiogr. 2011;24:60–74.
- 14. Olivieri L, Arling B, Friberg M, Sable C. Coronary artery Z score regression equations and calculators derived from a large heterogeneous population of children undergoing echocardiography. J Am Soc Echocardiogr. 2009;22:159–64.

- 15. Manlhiot C, Millar K, Golding F, McCrindle BW. Improved classification of coronary artery abnormalities based only on coronary artery z-scores after Kawasaki disease. Pediatr Cardiol. 2010; 31:242–9.
- 16. Tan TH, Wong KY, Cheng TK, Heng JT. Coronary normograms and the coronary-aorta index: objective determinants of coronary artery dilatation. Pediatr Cardiol. 2003;24:328–35.
- 17. Dominiguez SR, Anderson MS, El-Adawy M, Glodé MP. Preventing coronary artery abnormalities: a need for earlier diagnosis and treatment for Kawasaki disease. Pediatr Infect Dis J. 2012;31:1217–20.
- 18. Cameron SA, Carr M, Pahl E, DeMarais N, Shulman ST, Rowley AH. Coronary artery aneurysms are more severe in infants than in older children with Kawasaki disease. Arch Dis Child. 2019;104:451–5.
- 19. Mastrangelo G, Cimaz R, Calabri GB, Sinonimi G, Lasagni D, Resti M, et al. Kawasaki disease in infants less than one year of age: an Italian cohort from a single center. BMC Pediatrics. 2019;19:321.
- 20. Takahashi K, Oharaseki T, Yokouchi Y. Histopathological aspects of cardiovascular lesions in Kawasaki disease. Int J Rheum Dis. 2018;21:31–5.
- 21. Orenstein JM, Shulman ST, Fox LM, Baker SC, Takahashi M, Bhatti TR, et al. Three linked vasculopathic processes characterize Kawasaki disease: a light and transmission electron microscopic study. PLoS One. 2012;7:e38998.
- 22. Hörl M, Michel H, Döring S, Dechant MJ, Zeman F, Melter M, et al. Value of serial echocardiography in diagnosing Kawasaki's disease. Eur J Pediatr. 2021;180:387–95.
- 23. Noto N, Komori A, Ayusawa M, Takahashi S. Recent updates on echocardiography and ultrasound for Kawasaki disease: beyond the coronary artery. Cardiovasc Diagn Ther. 2018;8:80–9.
- 24. de Zorzi A, Colan SD, Gauvreau K, Baker AL, Sundel RP, Newburger JW. Coronary artery dimensions may be misclassified as normal in Kawasaki disease. J Pediatr. 1998;133:254–8.
- 25. Krzanowski M, Bodzoń W, Dimitrow PP. Imaging of all three coronary arteries by transthoracic echocardiography. an illustrated guide. Cardiovasc Ultrasound. 2003;1:16.
- 26. Brown LM, Duffy CE, Mitchell C, Young L. A practical guide to pediatric coronary artery imaging with echocardiography. J Am Soc Echocardiogr. 2015;28:379–91.
- 27. Muniz JC, Dummer K, Gauvreau K, Colan SD, Fulton DR, Newburger JW. Coronary artery dimensions in febrile children without Kawasaki disease. Circ Cardiovasc Imaging. 2013;6:239–44.
- 28. JCS Joint working Group. Guidelines for diagnosis and management of cardiovascular sequelae in Kawasaki disease (JCS 2008)—digest version. Circ J. 2010;74:1989–2020.
- 29. Ravekes WJ, Colan SD, Gauvreau K, Baker AL, Sundel RP, van der Velde ME, et al. Aortic root dilation in Kawasaki disease. Am J Cardiol. 2001;87:919–22.
- 30. Printz BF, Sleeper LA, Newburger JW, Minich LL, Bradley T, Cohen MS, et al.; Pediatric Heart Network Investigators. Noncoronary cardiac abnormalities are associated with coronary artery dilation and with laboratory inflammatory markers in acute Kawasaki disease. J Am Coll Cardiol. 2011:57:86–92.
- 31. Tsuchihashi T, Kakimoto N, Takeuchi T, Suenaga T, Suzuki T, Shibuta S, et al. Intimal thickening and disruption of the media occur in the arterial walls of coronary arteries not associated with coronary arterial aneurysms in patients with Kawasaki disease. BMC Cardiovasc Disord. 2021;21:278.
- 32. Dionne A, Ibrahim R, Gebhard C, Bakloul M, Selly JB, Leye M, et al. Coronary wall structural changes in patients with Kawasaki disease: new insights from optical coherence tomography (OCT). J Am Heart Assoc. 2015;4:e001939.